arXiv:math/0009141v1 [math.QA] 14 Sep 2000

Heisenberg double, pentagon equation, structure and classification of finite dimensional Hopf algebras

G. Militaru
University of Bucharest
Faculty of Mathematics
Str. Academiei 14
RO-70109 Bucharest 1, Romania
email: gmilit@al.math.unibuc.ro

Abstract

The study at the lavel of algebras of the pentagon equation $R^{12}R^{13}R^{23} = R^{23}R^{12}$ leds to the Structure and Classification theorem for finite dimensional Hopf algebras. We shall prove that L is a finite dimensional Hopf algebra if and only if there exists an invertible matrix $R = (A_{ij})_{i,j=1,n} \in \mathcal{M}_{n^2}(k) \cong \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$, solution of the pentagon equation $\sum_{j=1}^{n} A_{ij} \otimes A_{jp} = R(A_{ip} \otimes I_n)R^{-1}$, such that $L \cong P(\mathcal{M}_n(k), R) = P(n, R)$. Finally, we prove the Classification Theorem: there exists a one to one correspondence between the set of types of n-dimensional Hopf algebras and the set of orbits of the action $GL_n(k) \times (\mathcal{M}_n(k) \otimes \mathcal{M}_n(k)) \to \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$, $(u, R) \to (u \otimes u)R(u \otimes u)^{-1}$, the representatives of which are invertible solutions of length n for the pentagon equation.

Introduction

The theory of Hopf algebras proved to be a unifying framework for problems arising from different fields like: groups theory, Galois theory, affine algebraic groups, Lie algebras, local compact groups, operator algebras and quantum mechanics. Recently, using the classification of finite groups as a model, first steps were taken towards the classification of certain types of finite dimensional Hopf algebras. The techniques used are extremely diverse and the theory of Hopf algebras classification is far from being completed (see the review paper [13] and its list of references).

In this paper we shall study, at the lavel of algebras, the pentagon equation (also called the fusion equation) $R^{12}R^{13}R^{23} = R^{23}R^{12}$, untill now analyzed mainly in the theory of C^*-algebras (see [2] and the references indicated here). Let G be a locally compact group and dg a right Haar measure on G. Then the operator $V_G(\xi)(s,t) = \xi(st, t)$ is a solution of the pentagon equation. Let H be a separable Hilbert space and $R \in \mathcal{L}(H \otimes H)$ be a unitary solution of the pentagon equation. If R is commutative, then R is equivalent to

$V_G \otimes \mathrm{Id}_{K\otimes K}$, for a locally compact group G and a Hilbert space K [2, Theorem 2.2]. If H is finite dimensional, then [2, Theorem 4.7 and Theorem 4.10] proves that R is equivalent to $V_S \otimes \mathrm{Id}_{K\otimes K}$ where, this time, S is the Woronowicz C^*-algebra associated to R. R. M. Kashaev shows in [8] that the operator V_G is in fact the "coordinate" representation of the canonical element associated to the Heisenberg double of the group algebra kG. This remark will play an important role in proving the structure of $\underline{Pent}$, the category of pentagon objects: the objects of it are pairs (A, R) where A is a finite dimensional algebra and $R \in A \otimes A$ is an invertible solution of the pentagon equation $R^{12}R^{13}R^{23} = R^{23}R^{12}$ in $A \otimes A \otimes A$. Let $\underline{Hopf}$ be the category of finite dimensional Hopf algebras, $L \in \underline{Hopf}$ and $\mathcal{H}(L) = L\#L^*$ the Heisenberg double of L. $\mathcal{H}(L)$ is an algebra (we shall prove, using Morita theory, that it is isomorphic to the matrix algebra $\mathcal{M}_{\dim(L)}(k)$) the representations of which are the classic Hopf modules ${}^L\mathcal{M}_L$. [17, Theorem 5.2] and [8, Theorem 1] proves that the canonical element $\mathcal{R} \in \mathcal{H}(L) \otimes \mathcal{H}(L)$ of the Heisenberg double is a solution of the pentagon equation. Using a similar construction to the one of P. S. Baaj and G. Skandalis ([2]), we shall prove that there exists a functor

$$P : \underline{Pent} \to \underline{Hopf}, \quad (A, R) \to P(A, R)$$

such that $L \cong P(\mathcal{H}(L), \mathcal{R})$ for any $L \in \underline{Hopf}$. The structure of the solutions for the pentagon equation shows that, up to an algebra map, these solutions are canonical elements of varoius Heisebergs: i.e. $(A, R) \in \underline{Pent}$ if and only if there exists $L \in \underline{Hopf}$ and an algebra map $F : \mathcal{H}(L) \to A$ such that $R = (F \otimes F)(\mathcal{R})$, where $\mathcal{R} \in \mathcal{H}(L) \otimes \mathcal{H}(L)$ is the canonical element of the Heisenberg double $\mathcal{H}(L)$. The role of the Hopf modules category in solving the pentagon equation was evidenced before in [15, Proposition 1.3] and independently in [12], where, in the equivalent context of the Hopf equation, a FRT type theorem was proved. Using the Fundamental Theorem for Hopf modules, a Lagrange type theorem is proven: for $(A, R) \in \underline{Pent}$, A is free as a right $P(A, R)$-module and $\dim(A) = \dim(P(A, R))\dim(A^{R,r})$, where $A^{R,r}$ is the subspace of what we called the right R-coinvariants of A. The most important consequence of the results mentionated above is the fact that they led us to the Structure and the Classification theorem for finite dimensional Hopf algebras. Theorem 3.2 proves that L is a finite dimensional Hopf algebra if and only if there exists a invertible matrix $R \in \mathcal{M}_n(k) \otimes \mathcal{M}_n(k) \cong \mathcal{M}_{n^2}(k)$, solution of the pentagon equation, such that $L \cong P(\mathcal{M}_n(k), R) = P(n, k)$. The actual description, in terms of R, of the Hopf algebra structure of $P(n, R)$ is obviuosly included: $P(n, R)$ is the subalgebra of $\mathcal{M}_n(k)$ of the left coefficients of R with the comultiplication given by $\Delta(x) = R^{-1}(I_n \otimes x)R$, for all $x \in P(n, R)$. By writing R as $R = (A_{ij})_{i,j=1,n} \in \mathcal{M}_{n^2}(k) \cong \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$, for some matrices $A_{ij} \in \mathcal{M}_n(k)$, we get a much more elegant form of the pentagon equation for R

$$\sum_{j=1}^{n} A_{ij} \otimes A_{jp} = R(A_{ip} \otimes I_n)R^{-1}$$

for all i, $p = 1, \cdots n$.[1] Theorem 3.3 is the Classification Theorem: there exists a one to one correspondence between the set of types of n-dimensional Hopf algebras and the set of orbits of the action

$$GL_n(k) \times (\mathcal{M}_n(k) \otimes \mathcal{M}_n(k)) \to \mathcal{M}_n(k) \otimes \mathcal{M}_n(k), \quad (u, R) \to (u \otimes u)R(u \otimes u)^{-1},$$

---

[1] If A and $B \in \mathcal{M}_n(k)$, $A \otimes B \in \mathcal{M}_{n^2}(k)$ is the Kronecker product of A and B.

the representatives of which are invertible solutions of length n for the pentagon equation. In this way, the theory of finite dimensional Hopf algebras classification is reduced to a new Jordan type theory, which we called *restricted Jordan theory*, to be studied in more depth in another paper.

1 Preliminaries

Throughout this paper, k will be a commutative field. Unless specified otherwise, all vector spaces, algebras, Hopf algebras, tensor products and homomorphisms are over k. Let V be a finite dimensional vector space and $\varphi : V\otimes V^* \to \mathrm{End\,(V)}$, $\varphi(v\otimes v^*)(w) = \langle v^*, w\rangle v$ be the canonical isomorphism. We recall that the element $R := \varphi^{-1}(\mathrm{Id}_V)$ is called the *canonical element* of $V\otimes V^*$. Of course

$$R = \sum_{i=1}^{n} e_i \otimes e_i^*$$

where $\{e_i, e_i^* \mid i = 1, \cdots, n\}$ is a dual basis and R is independent of the choice of the dual basis.

For a k-algebra A, $\mathcal{M}_A$ (resp. ${}_A\mathcal{M}$) will be the category of right (resp. left) A-modules and A-linear maps. $\mathcal{M}_n(k)$ will be the k-algebra of $n\times n$-matrices and $GL_n(k)$ will be the group of invertible $n\times n$-matrices. We shall denote by (e_{ij}) the canonical basis of $\mathcal{M}_n(k)$: e_{ij} is the matrix having 1 in the (i,j)-position and 0 elsewhere. An n-dimensional k-algebra A is viewed as a subalgebra of $\mathcal{M}_n(k)$, via: $A\subset \mathrm{End\,(A)} \cong \mathcal{M}_\mathrm{n}(\mathrm{k})$. If $A = (a_{ij})$, $B = (b_{ij}) \in \mathcal{M}_n(k)$ then, via the canonical isomorphism $\mathcal{M}_n(k)\otimes\mathcal{M}_n(k) \cong \mathcal{M}_{n^2}(k)$, $A\otimes B$ viewed as a matrix of $\mathcal{M}_{n^2}(k)$ is given by the Kronecker product:

$$A\otimes B = \begin{pmatrix} a_{11}B & \cdots & a_{1n}B \\ \cdot & \cdots & \cdot \\ a_{n1}B & \cdots & a_{nn}B \end{pmatrix} \tag{1}$$

Let A be a k-algebra and $0 \neq R = \sum R^1\otimes R^2 \in A\otimes A$. Let $R_{(l)}$, $R_{(r)}$ be the subspaces of left, respectively right coefficients of R, i.e.

$$R_{(l)} = \{\sum \langle a^*, R^2\rangle R^1 \mid a^*\in A^*\}, \quad R_{(r)} = \{\sum \langle a^*, R^1\rangle R^2 \mid a^*\in A^*\}$$

Let us assume that $R = \sum_{i=1}^m a_i\otimes b_i$, where m is as small as possible. Then m is called the *length* of R and will be denoted $l(R) = m$. From the choice of m, the sets $\{a_i \mid i = 1, \cdots, m\}$, respectively $\{b_i \mid i = 1, \cdots, m\}$ are linear independent in A and hence bases of $R_{(l)}$, respectively $R_{(r)}$. In particular, $\dim(R_{(l)})$=$\dim(R_{(r)})$=$l(R)$. Two elements R and $S\in A\otimes A$ are called *equivalent* (we shall write $R\sim S$) if there exists $u\in U(A)$ an invertible element of A such that $S = {}^uR := (u\otimes u)R(u\otimes u)^{-1}$. If $R\sim S$ then $l(R)$=$l(S)$. In particular, if $\{a_i \mid i = 1, \cdots, m\}$ is a basis of $R_{(l)}$, then $\{ua_iu^{-1} \mid i = 1, \cdots, m\}$ is a basis of ${}^uR_{(l)}$.

For coproducts and comodules we use Sweedler's notation with suppressed summation sign: if L is a Hopf algebra, then for all $l\in L$ we write $\Delta(l) = l_{(1)}\otimes l_{(2)}$, $(\Delta\otimes \mathrm{Id})\Delta(l) = (\mathrm{Id}\otimes\Delta)\Delta(l) = l_{(1)}\otimes l_{(2)}\otimes l_{(3)}$. If (M, ρ_M) is a left L-comodule, then we write $\rho_M(m) = m_{<-1>}\otimes m_{<0>} \in L\otimes M$, for all $m\in M$. ${}^L\mathcal{M}$ (resp. $\mathcal{M}^L$) will be the category of left (resp. right) L-comodules and L-colinear maps.

1.1 The Heisenberg double and Hopf modules

Let L be a Hopf algebra with a bijective antipode. We recall that a right-left L-Hopf module is a threetuple $(M, \cdot, \rho)$, where $(M, \cdot)$ is a right L-module, (M, ρ) is a left L-comodule such that the following compatibility relation holds

$$\rho(m \cdot l) = \rho(m)\Delta(l) \tag{2}$$

for all $l \in L$, $m \in M$. ${}^{L}\mathcal{M}_L$ will be the category of right-left L-Hopf modules and L-linear, L-colinear maps. If V is a vector space, then $V \otimes L$ is a L-Hopf module via $(v \otimes l) \cdot g := v \otimes lg$, and $v \otimes l \to l_{(1)} \otimes v \otimes l_{(2)}$, for all $v \in V$, l, $g \in L$. Thus we have a functor $- \otimes L : \mathcal{M}_k \to {}^{L}\mathcal{M}_L$, $V \to V \otimes L$. Let $(-)^{\mathrm{co}(L)} : {}^{L}\mathcal{M}_L \to \mathcal{M}_k$, $M \to M^{\mathrm{co}(L)} = \{m \in M \mid \rho(m) = 1_L \otimes m\}$ be the functor that associates to each M its coinvariants. The Fundamental Theorem for Hopf modules [16] (in the right-left version) shows that for any L-Hopf module M the multiplication map $\mu : M^{\mathrm{co}(L)} \otimes L \to M$, $\mu(m \otimes l) = m \cdot l$ is an isomorphism of L-Hopf modules with the inverse $\beta : M \to M^{\mathrm{co}(L)} \otimes L$, $\beta(m) = m_{<0>} \cdot S^{-1}(m_{<-1>}) \otimes m_{<-2>}$. Hence, $(- \otimes L) \circ (-)^{\mathrm{co}(L)} \cong \mathrm{Id}_{{}^{L}\mathcal{M}_L}$. As we also have that $(-)^{\mathrm{co}(L)} \circ (- \otimes L) \cong \mathrm{Id}_{\mathcal{M}_k}$, we obtain that the functors $- \otimes L : \mathcal{M}_k \to {}^{L}\mathcal{M}_L$, $(-)^{\mathrm{co}(L)} : {}^{L}\mathcal{M}_L \to \mathcal{M}_k$ define an equivalence of categories. The same is valid for all left-right types of L-Hopf modules: ${}_{L}\mathcal{M}^{L}$, $\mathcal{M}_L^L$ or ${}_{L}^{L}\mathcal{M}$. For instance, let ${}_{L}\mathcal{M}^{L}$ be the category of left-right L-Hopf modules whose objects are left L-modules and right L-comodules M such that $\rho(l \cdot m) = \Delta(l)\rho(m)$ for all $l \in L$, $m \in M$. For this category the inverse of the multiplication map $L \otimes M^{\mathrm{co}(L)} \to M$, in the Fundamental Theorem, is given by $\alpha(m) = m_{<2>} \otimes S^{-1}(m_{<1>}) \cdot m_{<0>}$, for all $m \in M$. To conclude, if the antipode of L is bijective, then we have the equivalences of categories

$${}^{L}\mathcal{M}_L \cong \mathcal{M}_k \cong {}_{L}\mathcal{M}^{L}.$$

We recall that L^* is a left L-module algebra in the usual way $\langle h \cdot g^*, h' \rangle = \langle g^*, h'h \rangle$ for all h, $h' \in L$, $g^* \in L^*$. The *Heisenberg double* of L is by definition the smash product

$$\mathcal{H}(L) = L\#L^*,$$

i.e. $\mathcal{H}(L) = L \otimes L^*$ as a vector space, and the multiplication is given by

$$(h\#h^*)(g\#g^*) = h_{(2)}g\#h^*(h_{(1)} \cdot g^*) = h_{(2)}g\#h^*\langle g^*, ?h_{(1)} \rangle$$

for all h, $g \in L$, h^*, $g^* \in L^*$. $\mathcal{H}(L)$ is an associative algebra with the unit $1_L\#\varepsilon_L$. The Heisenberg double $\mathcal{H}(L) = L\#L^*$ is in fact a particular case (for the Doi-Koppinen datum $(H, A, C) = (L, L, L)$) of the general smash product $A\#C^*$ introduced in [6] in the right-right version. The right-left version of it and the above description of the Heisenberg double is given in [4]. The canonical isomorphism of vector spaces

$$\varphi : \mathcal{H}(L) \to \mathrm{End\,(L)} = \mathcal{M}_{\mathrm{dim(L)}}(\mathrm{k}), \quad \varphi(\mathrm{l}\#\mathrm{l}^*)(\mathrm{g}) = \langle \mathrm{l}^*, \mathrm{g} \rangle \mathrm{l}$$

is not an algebra map. However, we can prove the following

Proposition 1.1 *Let L be a finite dimensional Hopf algebra. Then there exists an algebra isomorphism*

$$\mathcal{H}(L) \cong \mathcal{M}_{\dim(L)}(k).$$

Proof. As L is finite dimensional, the functor

$$T : {}^{L}\mathcal{M}_L \to \mathcal{M}_{\mathcal{H}(L)}, \quad T(M) = M$$

where the right $\mathcal{H}(L)$-action on M is given by

$$m \bullet (l\#l^*) = \langle l^*, m_{<-1>}\rangle m_{<0>} \cdot l$$

is an equivalence of categories ([6]). As the antipode of L is bijective ([16]) we have the following equivalences of categories

$$\mathcal{M}_{\mathcal{H}(L)} \cong {}^{L}\mathcal{M}_L \cong \mathcal{M}_k$$

i.e. $\mathcal{H}(L)$ is Morita equivalent to k. It follows from the Morita theory that there exists an algebra isomorphism $\mathcal{H}(L) \cong \mathcal{M}_n(k)$. Taking into account that $\dim(\mathcal{H}(L)) = \dim(L)^2$, we obtain that $n = \dim(L)$. □

Remark 1.2 The k-linear maps

$$i_L : L \to \mathcal{H}(L),\ i_L(l) = l\#\varepsilon_L \quad \text{and} \quad i_{L^*} : L^* \to \mathcal{H}(L),\ i_{L^*}(l^*) = 1_L\#l^*$$

for all $l \in L$, $l^* \in L^*$ are injective algebra maps. Beeing a smash product, the Heisenberg double $\mathcal{H}(L)$ satisfies the following universal property: given an algebra A and the algebra maps $u : L \to A$, $v : L^* \to A$ such that

$$u(l)v(l^*) = v(l_{(1)} \cdot l^*)u(l_{(2)}) \tag{3}$$

there exists a unique algebra map $F : \mathcal{H}(L) \to A$ (given by $F(l\#l^*) := v(l^*)u(l)$, for all $l \in L$, $l^* \in L^*$) such that

$$F \circ i_L = u, \quad F \circ i_{L^*} = v \tag{4}$$

1.2 The pentagon equation

Definition 1.3 *Let A be an algebra and $R = \sum R^1 \otimes R^2 \in A \otimes A$ be an invertible element. We shall say that R is a solution of the pentagon equation if*

$$R^{12}R^{13}R^{23} = R^{23}R^{12} \tag{5}$$

in $A \otimes A \otimes A$. R is called unitary if $1_A \in R_{(l)} \cap R_{(r)}$, i.e. there exist λ, $\omega \in A^*$ such that

$$\sum \langle \lambda, R^1\rangle R^2 = \sum \langle \omega, R^2\rangle R^1 = 1_A \tag{6}$$

Pent will be the category of the (finite) pentagon objects: the objects of it are pairs (A, R), where A is a finite dimensional algebra and $R \in A \otimes A$ is an invertible solution of the pentagon equation. A morphism $f : (A, R) \to (B, T)$ between two pentagon objects (A, R) and (B, T) is an algebra map $f : A \to B$ such that $(f \otimes f)(R) = T$. Pent is a monoidal category under the product $(A, R) \otimes (B, T) := (A \otimes B, R^{13}T^{24})$.

Let $R \in A \otimes A$ be an invertible solution of the pentagon equation. The subspaces of A

$$A^{R,l} = \{a \in A \mid R(a \otimes 1_A) = a \otimes 1_A\} \quad \text{and} \quad A^{R,r} = \{a \in A \mid (1_A \otimes a)R = 1_A \otimes a\}$$

are called the spaces of left, respectively right, R-*coinvariants* of A. On the other hand, the subspaces

$$^{R,l}A = \{a \in A \mid R(1_A \otimes a) = 1_A \otimes a\} \quad \text{and} \quad ^{R,r}A = \{a \in A \mid (a \otimes 1_A)R = a \otimes 1_A\}$$

are called the spaces of left, respectively right, R-*invariants* of A.

Remarks 1.4 1. R is a solution for the pentagon equation if and only if $T = R^{-1}$ is a solution of what we called in [12] the *Hopf equation*: $T^{12}T^{23} = T^{23}T^{13}T^{12}$.
2. Let $R \in A \otimes A$ be an invertible solution of the pentagon equation. If the subspaces $A^{R,l}$ and $A^{R,r}$ are non-zero then R is unitary. Indeed, let $a \neq 0$ be an element of $A^{R,l}$ and $R = \sum_{i=1}^{m} a_i \otimes b_i \in A \otimes A$, where $m = l(R)$. Then $\sum_{i=1}^{m} a_i a \otimes b_i = a \otimes 1_A$ and hence, $1_A = \sum_{i=1}^{m} \langle a^*, a_i a\rangle b_i \in R_{(r)}$, where $a^* \in A^*$ such that $\langle a^*, a\rangle = 1$. Similarly, if $A^{R,r} \neq 0$ then $1_A \in R_{(l)}$. It follows from Corollary 2.2 that an invertible solution of the pentagon equation is unitary if and only if the spaces $A^{R,l}$ and $A^{R,r}$ are nonzero. On the other hand, Corollary 2.2 proves that if A is finite dimensional, any invertible solution of the pentagon equation is unitary.
3. In some applications, the algebra A is of the form $A = \mathrm{End}_k(M)$, where M is a vector space over k. Then we can view R as an element of $\mathrm{End}_k(M \otimes M)$, using the natural embedding $\mathrm{End}_k(M) \otimes \mathrm{End}_k(M) \subseteq \mathrm{End}_k(M \otimes M)$, and the pentagon equation as an equation in $\mathrm{End}_k(M \otimes M \otimes M)$. It was proved in [5, Proposition 7.1] that there exists a bijective correspondence between all structures of monoidal categories on the category of vector spaces over k and all bijective solutions $R \in \mathrm{End}_k(M \otimes M)$ of the pentagon equation on various k-vector spaces M.

Examples 1.5 1. Let $R \in A \otimes A$ be an invertible solution of the pentagon equation and $f : A \to B$ an algebra map. Then $(f \otimes f)(R) \in B \otimes B$ is an invertible solution of the pentagon equation.
2. If $R \in A \otimes A$ is a unitary invertible solution of the pentagon equation and $u \in U(A)$ is an invertible element of A, then $^{u}R = (u \otimes u)R(u \otimes u)^{-1}$ is a unitary invertible solution of the pentagon equation.
3. Let A be an algebra a, $b \in A$ and $R \in A \otimes A$ given by

$$R = 1_A \otimes 1_A + a \otimes b.$$

Then R is a solution of the pentagon equation if and only if

$$a \otimes (ab - ba - 1) \otimes b = a \otimes a \otimes b^2 + a^2 \otimes b \otimes b + a^2 \otimes ba \otimes b^2. \tag{7}$$

Now, if we take a and b such that

$$a^2 = b^2 = 0 \quad \text{and} \quad ab - ba = 1 \tag{8}$$

or

$$a^2 = 0, \quad b^2 = b \quad \text{and} \quad ab - ba = a + 1 \tag{9}$$

then (7) is satisfied and R is an invertible solution of the pentagon equation.
We shall give some specific examples of these types in the case that $A = \mathcal{M}_n(k)$. First, in both cases, it follows from the last equation that $\mathrm{char}(k)|n$.
Assume now that $\mathrm{char}(k) = 2$ and let $n = 2q$, where q is a positive integer. Then

$$a = e_{12} + e_{34} + \cdots + e_{2q-1,2q}, \quad b = e_{21} + e_{43} + \cdots + e_{2q,2q-1}$$

is a solution of (8), and hence $R = I_n \otimes I_n + \sum_{i,j=1}^{q} e_{2i-1,2i} \otimes e_{2j,2j-1}$ is an invertible solution of the pentagon equation. On the other hand, for an arbitrary invertible matrix $X \in \mathcal{M}_q(k)$, the matrices a, $b \in \mathcal{M}_n(k)$ given by

$$b = e_{11} + e_{22} + \cdots e_{qq}, \quad a = \begin{pmatrix} I_q & X^{-1} \\ X & I_q \end{pmatrix}$$

are solutions of (9) and hence $R = R_X = I_n \otimes I_n + \sum_{i=1}^{q} a \otimes e_{ii}$ is an invertible solution of the pentagon equation.
4. Let $(M, \cdot, \rho) \in {}_L\mathcal{M}^L$ be a L-Hopf module over a Hopf algebra L. Then the map

$$R_{(M,\cdot,\rho)} : M \otimes M \to M \otimes M, \quad R_{(M,\cdot,\rho)}(m \otimes n) = m_{<0>} \otimes m_{<1>} \cdot n \tag{10}$$

is a bijective solution of the pentagon equation in $\mathrm{End}_{\,\mathrm{k}}(\mathrm{M}\otimes\mathrm{M}\otimes\mathrm{M})$ ([12], [15]). Conversely, if M is finite dimensional and $R \in \mathrm{End}_{\,\mathrm{k}}(\mathrm{M}\otimes\mathrm{M})$ is a solution of the pentagon equation, then there exists a bialgebra $B(R)$ such that $(M, \cdot, \rho) \in {}_{B(R)}\mathcal{M}^{B(R)}$ and $R = R_{(M,\cdot,\rho)}$; this is [12, Theorem 3.1] taking into account the equivalence between the Hopf and the pentagon equation: R is a solution of the pentagon equation in $\mathrm{End}_{\,\mathrm{k}}(\mathrm{M}\otimes\mathrm{M}\otimes\mathrm{M})$ if and only if $\tau R \tau$ is a solution of the Hopf equation, where $\tau \in \mathrm{End}_{\,\mathrm{k}}(\mathrm{M}\otimes\mathrm{M})$ is the flip map.

Proposition 1.6 *Let A be an algebra and $R \in A \otimes A$ an invertible solution of the pentagon equation. Consider the comultiplications Δ_r, $\Delta_l : A \to A \otimes A$, given by*

$$\Delta_r(a) = R^{-1}(1_A \otimes a)R = \sum U^1R^1 \otimes U^2aR^2 \tag{11}$$

$$\Delta_l(a) = R(a \otimes 1_A)R^{-1} = \sum R^1aU^1 \otimes R^2U^2 \tag{12}$$

where $U = \sum U^1 \otimes U^2 = R^{-1}$. Then $A_r = (A, \cdot, \Delta_r)$ and $A_l = (A, \cdot, \Delta_l)$ are bialgebras without counit.

Proof. It is obvious that Δ_r, Δ_l are algebra maps. For $a \in A$ we have

$$(\mathrm{Id} \otimes \Delta_r)\Delta_r(a) = (R^{23})^{-1}(R^{13})^{-1}(1_A \otimes 1_A \otimes a)R^{13}R^{23}$$

and

$$(\Delta_r \otimes \mathrm{Id})\Delta_r(a) = (R^{12})^{-1}(R^{23})^{-1}(1_A \otimes 1_A \otimes a)R^{23}R^{12}$$

so Δ_r is coassociative if and only if

$$R^{23}R^{12}(R^{23})^{-1}(R^{13})^{-1}(1_A \otimes 1_A \otimes a) = (1_A \otimes 1_A \otimes a)R^{23}R^{12}(R^{23})^{-1}(R^{13})^{-1}.$$

Using the pentagon equation (5), we find that this is equivalent to

$$R^{12}(1_A \otimes 1_A \otimes a) = (1_A \otimes 1_A \otimes a)R^{12}$$

and this equality holds for any $a \in A$. In a similar way we can prove that Δ_l is also coassociative. □

It follows from Proposition 1.6 that we can put two different algebra structures (without unit) on the dual A^*: the multiplications are the convolutions $*_l$ and $*_r$ which are the dual maps of Δ_l and Δ_r, i.e.

$$\langle \omega *_l \omega', a\rangle = \sum \langle \omega, R^1 a U^1\rangle \langle \omega', R^2 U^2\rangle \tag{13}$$

$$\langle \omega *_r \omega', a\rangle = \sum \langle \omega, U^1 R^1\rangle \langle \omega', U^2 a R^2\rangle \tag{14}$$

for all ω, $\omega' \in A^*$, $a \in A$.
Aside from the invertibility condition for $\mathcal{R}$, the following theorem is [17, Theorem 5.2] and [8, Theorem 1]. In [17], the Heisenberg double does not appear explicitely, and in [8] the Heisenberg double is described in terms of structure constants, and not as a smash product. For this reason, and for the reader's convenience, we shall present the proof.

Theorem 1.7 *Let L be a finite dimensional Hopf algebra and $\{e_i, e_i^* \mid i = 1, \cdots, n\}$ a dual basis of L. Then the canonical element*

$$\mathcal{R} = \sum_i (e_i \# \varepsilon) \otimes (1 \# e_i^*) \in \mathcal{H}(L) \otimes \mathcal{H}(L)$$

is an invertible solution of the pentagon equation in $\mathcal{H}(L) \otimes \mathcal{H}(L) \otimes \mathcal{H}(L)$.
Consequently, if A is an algebra and $f : \mathcal{H}(L) \to A$ an algebra map, then $(f \otimes f)(\mathcal{R})$ is an invertible solution of the pentagon equation in $A \otimes A \otimes A$.

Proof. Taking into account the multiplication rule of $\mathcal{H}(L)$ we have

$$\mathcal{R}^{23}\mathcal{R}^{12} = \sum_{i,j} (e_j \# \varepsilon) \otimes (e_{i_{(2)}} \# e_{i_{(1)}} \cdot e_j^*) \otimes (1 \# e_i^*)$$

and

$$\mathcal{R}^{12}\mathcal{R}^{13}\mathcal{R}^{23} = \sum_{a,b,c} (e_a e_b \# \varepsilon) \otimes (e_c \# e_a^*) \otimes (1 \# e_b^* e_c^*)$$

so we have to prove the equality

$$\sum_{i,j} e_j \otimes e_{i_{(2)}} \otimes e_{i_{(1)}} \cdot e_j^* \otimes e_i^* = \sum_{a,b,c} e_a e_b \otimes e_c \otimes e_a^* \otimes e_b^* e_c^* \tag{15}$$

in $L \otimes L \otimes L^* \otimes L^*$. Let's fix the indices x, y, z and $t = \{1, \cdots, n\}$ and evaluate (15) at $e_x^* \otimes e_y^* \otimes e_z \otimes e_t$. (15) is then equivalent to

$$\langle e_y^*, e_{t_{(2)}} \rangle \langle e_x^*, e_z e_{t_{(1)}} \rangle = \sum_b \langle e_x^*, e_z e_b \rangle \langle e_b^* e_y^*, e_t \rangle,$$

which can be obtained by applying the definition of the convolution product and the dual basis formula in the right hand side. We shall prove now that

$$U = \sum_i (S(e_i) \# \varepsilon) \otimes (1 \# e_i^*)$$

is the inverse of R, where S is the antipode of L. Since $\mathcal{H}(L) \otimes \mathcal{H}(L)$ is isomorphic to $\mathcal{M}_{n^2}(k)$, it is enough to prove that $RU = 1 \otimes 1$. As $RU = \sum_{i,j} (e_i S(e_j) \# \varepsilon) \otimes (1 \# e_i^* e_j^*)$, we have to prove the formula

$$\sum_{i,j} e_i S(e_j) \otimes e_i^* e_j^* = 1 \otimes \varepsilon$$

which holds, as for indices x, $y = 1, \cdots, n$ we have

$$\begin{aligned} \sum_{i,j} \langle e_x^*, e_i S(e_j) \rangle \langle e_i^* e_j^*, e_y \rangle &= \sum_{i,j} \langle e_x^*, e_i S(e_j) \rangle \langle e_i^*, e_{y_{(1)}} \rangle \langle e_j^*, e_{y_{(2)}} \rangle \\ &= \sum_{i,j} \langle e_x^*, e_i \langle e_i^*, e_{y_{(1)}} \rangle S(e_j \langle e_j^*, e_{y_{(2)}} \rangle) \rangle \\ &= \langle e_x^*, e_{y_{(1)}} S(e_{y_{(2)}}) \rangle = \langle e_x^*, 1 \rangle \langle \varepsilon, e_y \rangle \end{aligned}$$

□

2 The solutions of the Pentagon equation

We shall set some notations that will remain valid during the rest of the paper: $R = \sum R^1 \otimes R^2 \in A \otimes A$ will be an invertible solution of the pentagon equation, where A is an algebra. Let $R_{(l)}$, $R_{(r)}$ be the subspaces of left, respectively right coefficients of R. We shall denote them as follows

$$P = P(A, R) := R_{(l)}, \quad H = H(A, R) := R_{(r)}.$$

Assume now that $R = \sum_{i=1}^m a_i \otimes b_i$, where $m = l(R)$. The fact that m is minimal implies that the sets $\{a_i \mid i = 1, \cdots, m\}$ and $\{b_i \mid i = 1, \cdots, m\}$ are basis of $P = R_{(l)}$ and $H = R_{(r)}$. In particular, $\dim(P)$=$\dim(H) = l(R) = m$. Now consider $a_i^* \in P^*$ and $b_i^* \in H^*$ such that $\{a_i, a_i^*\}$ and $\{b_i, b_i^*\}$ are dual bases of P and H: $\langle a_i^*, a_j \rangle = \delta_{ij} = \langle b_i^*, b_j \rangle$. Extend $a_i^* : P \to k$ and $b_i^* : H \to k$ to respectively $\omega_i : A \to k$ and $\lambda_i : A \to k$. We then have

$$\sum_{i=1}^m \langle \omega_k, a_i \rangle b_i = b_k \quad \text{and} \quad \sum_{j=1}^m a_j \langle \lambda_k, b_j \rangle = a_k \tag{16}$$

for all $k = 1, \cdots, m$. We will use two different notations for R: $R = \sum_{i=1}^m a_i \otimes b_i = \sum_{j=1}^m a_j \otimes b_j$, when we are interested in the basis elements of P and H, and the generic notation $R = \sum R^1 \otimes R^2 = \sum r^1 \otimes r^2 = r$ and $U = R^{-1} = \sum U^1 \otimes U^2$.
The construction from part 1) of the next theorem is bassically the one of P. S. Baaj and G. Skandalis [2], for a unitary solution of the pentagon equation $R \in \mathcal{L}(H \otimes H)$, where H is a separable Hilbert space. Part 3) together with Theorem 1.7 give the structure of the category *Pent*. Part 4) is a Lagrange type theorem and the last part will play a role in the clasification of finite dimensional Hopf algebras.

Theorem 2.1 *Let A be an algebra, $R = \sum R^1 \otimes R^2 \in A \otimes A$ an unitary invertible solution of the pentagon equation and $P = P(A, R) = R_{(l)}$, $H = H(A, R) = R_{(r)}$ the subspaces of coefficients of R. Then:*
1) P and H are subalgebras of A and Hopf algebras with the comultiplications given by the formulas

$$\Delta_P : P \to P \otimes P, \quad \Delta_P(x) = \Delta_r(x) = R^{-1}(1_A \otimes x)R \tag{17}$$

$$\Delta_H : H \to H \otimes H, \quad \Delta_H(y) = \Delta_l(y) = R(y \otimes 1_A)R^{-1} \tag{18}$$

for all $x \in P$, $y \in H$. Furthermore, the subalgebra A_R of A generated by H and P is finite dimensional and $dim(A_R) \leq l(R)^2$.
2) The k-linear map $f : P^* \to H$, $f(p^*) = \sum \langle p^*, R^1 \rangle R^2$ is an isomorphism of Hopf algebras.
3) The k-linear map $F : \mathcal{H}(P) \to A$, $F(p\#p^*) = \sum \langle p^*, R^1 \rangle R^2 p$ is an algebra map and $R = (F \otimes F)(\mathcal{R})$, where $\mathcal{R} \in \mathcal{H}(P) \otimes \mathcal{H}(P)$ is the canonical element associated to the Heisenberg double.
4) The multiplication on A defines isomorphisms

$$A^{R,r} \otimes P \cong A \quad (\text{resp. } H \otimes A^{R,l} \cong A)$$

of right P-modules (resp. left H-modules). In particular, A is free as a right P-module and as a left H-module and, if A is finite dimensional

$$\dim(P) = \dim(H) = \frac{\dim(A)}{\dim(A^{R,l})} = \frac{\dim(A)}{\dim(A^{R,r})}.$$

5) If $f : A \to B$ is an algebra isomorphism and $S = (f \otimes f)(R)$, then the Hopf algebras $P(A, R)$ and $P(B, S)$ are isomorphic. Consequently, there exists an Hopf algebra isomorphism $P(A, R) \cong P(A, {}^uR)$, for any $u \in U(A)$.

Proof. 1) We shall use the notations introduced above. First we shall prove that P (resp. H) are unitary subalgebras in A and subcoalgebras of $A_r = (A, \Delta_r)$ (resp. $A_l = (A, \Delta_l)$). This will follow from the formulas:

$$a_p a_q = \sum_{j=1}^m \langle \lambda_p *_l \lambda_q, b_j \rangle a_j \in P, \quad \Delta_r(a_p) = \sum_{i,j=1}^m \langle \lambda_p, b_i b_j \rangle a_i \otimes a_j \in P \otimes P \tag{19}$$

and

$$b_p b_q = \sum_{j=1}^m \langle \omega_p *_r \omega_q, a_j \rangle b_j \in H, \quad \Delta_l(b_p) = \sum_{i,j=1}^m \langle \omega_p, a_i a_j \rangle b_i \otimes b_j \in H \otimes H \tag{20}$$

for all p, $q = 1, \cdots, m$. We prove (19), (being similar, (20) is left to the reader).

$$\begin{aligned}
\sum_{j=1}^{m} a_j \langle \lambda_p *_l \lambda_q, b_j \rangle &\overset{(13)}{=} \sum_{j=1}^{m} a_j \langle \lambda_p, R^1 b_j U^1 \rangle \langle \lambda_q, R^2 U^2 \rangle \\
&= (\mathrm{Id} \otimes \lambda_p \otimes \lambda_q)(\sum_{j=1}^{m} a_j \otimes R^1 b_j U^1 \otimes R^2 U^2 \rangle) \\
&= (\mathrm{Id} \otimes \lambda_p \otimes \lambda_q)(R^{23} R^{12} (R^{23})^{-1}) \overset{(5)}{=} (\mathrm{Id} \otimes \lambda_p \otimes \lambda_q)(R^{12} R^{13}) \\
&= (\mathrm{Id} \otimes \lambda_p \otimes \lambda_q)(\sum_{j,k=1}^{m} a_j a_k \otimes b_j \otimes b_k) \\
&= \sum_{j,k=1}^{m} a_j \langle \lambda_p, b_j \rangle a_k \langle \lambda_q, b_k \rangle = a_p a_q
\end{aligned}$$

i.e. P is a subalgebra of A. On the other hand

$$\begin{aligned}
\Delta_r(a_p) &\overset{(16)}{=} \Delta_r(\sum_{j=1}^{m} a_j \langle \lambda_p, b_j \rangle) \overset{(11)}{=} \sum_{j=1}^{m} U^1 R^1 \otimes U^2 a_j R^2 \langle \lambda_p, b_j \rangle \\
&= (\mathrm{Id} \otimes \mathrm{Id} \otimes \lambda_p)(\sum_{j=1}^{m} U^1 R^1 \otimes U^2 a_j R^2 \otimes b_j) \\
&= (\mathrm{Id} \otimes \mathrm{Id} \otimes \lambda_p)((R^{12})^{-1} R^{23} R^{12}) \overset{(5)}{=} (\mathrm{Id} \otimes \mathrm{Id} \otimes \lambda_p)(R^{12} R^{23}) \\
&= (\mathrm{Id} \otimes \mathrm{Id} \otimes \lambda_p)(\sum_{i,j=1}^{m} a_i \otimes a_j \otimes b_i b_j) = \sum_{i,j=1}^{m} a_i \otimes a_j \langle \lambda_p, b_i b_j \rangle
\end{aligned}$$

i.e. P is a subcoalgebra in (A, Δ_r). A similar computation yields (20), proving that H is a subalgebra of A and a subcoalgebra in (A, Δ_l). As R is unitary, $1_A \in P \cap H$, i.e. P and H are unitary subalgebras of A. Moreover, from the construction we have that $R \in P \otimes H \subset A \otimes A$ and hence we can view $U = R^{-1} \in P \otimes H$.
We shall define now the counit and the antipode of the Hopf algebras P and H. They are given by the formulas:

$$\varepsilon_P : P \to k, \quad \varepsilon_P(a_k) = \langle b_k^*, 1_A \rangle, \quad S_P : P \to P, \quad S_P(a_k) = \sum U^1 \langle b_k^*, U^2 \rangle \tag{21}$$

and

$$\varepsilon_H : H \to k, \quad \varepsilon_H(b_k) = \langle a_k^*, 1_A \rangle, \quad S_H : H \to H, \quad S_H(b_k) = \sum \langle a_k^*, U^1 \rangle U^2 \tag{22}$$

for all $k = 1, \cdots, m$. We shall prove that P is a Hopf algebra; the fact that H is a Hopf algebra is proved in a similar way. First, we remark that, as H is a subalgebra of A, the comultiplication from (19) can be rewritten as

$$\Delta_r(a_p) = \sum_{i,j=1}^{m} \langle b_p^*, b_i b_j \rangle a_i \otimes a_j \tag{23}$$

Now, for $p = 1, \cdots, m$ we have

$$\begin{aligned}(\mathrm{Id} \otimes \varepsilon_P)\Delta_r(a_p) &\overset{(23)}{=} \sum_{i,j=1}^{m} a_i \langle b_p^*, b_i b_j \rangle \langle b_j^*, 1_A \rangle = \sum_{i,j=1}^{m} a_i \langle b_p^*, b_i b_j \langle b_j^*, 1_A \rangle \rangle \\ &= \sum_{i=1}^{m} a_i \langle b_p^*, b_i \rangle = a_p\end{aligned}$$

i.e. $(\mathrm{Id} \otimes \varepsilon_P)\Delta_r = \mathrm{Id}$. A similar computation shows that $(\varepsilon_P \otimes \mathrm{Id})\Delta_r = \mathrm{Id}$, and ε_P is a counit of P. S_P is a right convolution inverse of Id_P since

$$\begin{aligned}(\mathrm{Id} \otimes S_P)\Delta_r(a_p) &\overset{(23)}{=} \sum_{i,j=1}^{n} a_i \langle b_p^*, b_i b_j \rangle U^1 \langle b_j^*, U^2 \rangle = \sum_{i,j=1}^{n} a_i U^1 \langle b_p^*, b_i b_j \langle b_j^*, U^2 \rangle \rangle \\ &= \sum_{i=1}^{n} a_i U^1 \langle b_p^*, b_i U^2 \rangle = (\mathrm{Id} \otimes b_p^*)(RR^{-1}) = 1_A \langle b_p^*, 1_A \rangle = \varepsilon_P(a_p) 1_A\end{aligned}$$

From the fact that P is finite dimensional, it follows that S_P is an antipode of P.
We shall prove now that the subalgebra A_R of A generated by H and P is finite dimensional. We shall use the pentagon equation. We have:

$$R^{12}R^{13}R^{23} = \sum_{i,j,k=1}^{m} a_i a_j \otimes b_i a_k \otimes b_j b_k \quad \text{and} \quad R^{23}R^{12} = \sum_{i,j=1}^{m} a_j \otimes a_i b_j \otimes b_i$$

Applying $a_u^* \otimes \mathrm{Id} \otimes b_v^*$ to the pentagon equation we obtain

$$a_u b_v = \sum_{i,j,k=1}^{m} \langle a_u^*, a_i a_j \rangle \langle b_v^*, b_j b_k \rangle b_i a_k$$

for all u, $v = 1, \cdots, m$. This formula gives that $\dim(A_R) \leq m^2$.
2) Let us prove now that $f : P^* \to H$ is an isomorphism of Hopf algebras. f is an isomorphism of vector spaces since $f(a_j^*) = \sum_{i=1}^{m} \langle a_j^*, a_i \rangle b_i = b_j$. The formula (20) can be rewritten as

$$b_p b_q = \sum_{j=1}^{m} \langle a_p^* *_r a_q^*, a_j \rangle b_j$$

which means that $f(a_p^*)f(a_q^*) = f(a_p^* *_r a_q^*)$ for all p, $q = 1, \cdots, m$, i.e. f is an algebra isomorphism. Let us prove now that f is also a coalgebra map. We recall the definition of the comultiplication Δ_{P^*}:

$$\Delta_{P^*}(a_p^*) = \sum X^1 \otimes X^2 \in P^* \otimes P^*, \text{ if and only if } \langle a_p^*, xy \rangle = \sum \langle X^1, x \rangle \langle X^2, y \rangle$$

for all x, $y \in P$. It follows that

$$\begin{aligned}(f \otimes f)\Delta_{P^*}(a_p^*) &= \sum f(X^1) \otimes f(X^2) = \sum \langle X^1, R^1 \rangle R^2 \otimes \langle X^2, r^1 \rangle r^2 \\ &= \sum \langle a_p^*, R^1 r^1 \rangle R^2 \otimes r^2 = \sum_{i,j=1}^{m} \langle a_p^*, a_i a_j \rangle b_i \otimes b_j \\ &\overset{(20)}{=} \Delta_H(b_p) = (\Delta_H \circ f)(a_p^*)\end{aligned}$$

i.e. f is also a coalgebra map. Hence, we have proved that f is an isomorphism of bialgebras and, as P and H are Hopf algebras, it is also a isomorphism of Hopf algebras ([16]).
3) We remark that

$$F(a_i \# a_j^*) = \sum_{t=1}^{m} \langle a_j^*, a_t \rangle b_t a_i = b_j a_i$$

for all i, $j = 1, \cdots, m$. The fact that F is an algebra map can be proved directly by using this formula; another way to proceed is to use the universal property of the Heisenberg double $\mathcal{H}(P)$ for the diagram (4), with $L = P$, $u : P \to A$ is the usual inclusion and $v : P^* \to A$ is the composition $v = f \circ j$, where $f : P^* \to H$ is the isomorphism from part 2) and $j : H \to A$ is the usual inclusion. We only have to prove that the compatibility condition (3) holds, i.e.

$$hv(g^*) = v(h_{(1)} \cdot g^*) h_{(2)}$$

for any $h \in P$ and $g^* \in P^*$, which turns out to be

$$\sum h \langle g^*, R^1 \rangle R^2 = \sum \langle g^*, R^1 h_{(1)} \rangle R^2 h_{(2)}$$

or, equivalently

$$\sum R^1 \otimes hR^2 = \sum R^1 h_{(1)} \otimes R^2 h_{(2)}.$$

This equation holds, as $\Delta_P(h) = R^{-1}(1_H \otimes h)R$, for any $h \in P$.
Now let $\mathcal{R} = \sum_{i=1}^{m} (a_i \# \varepsilon_P) \otimes (1_A \# a_i^*)$ be the canonical element of $\mathcal{H}(P) \otimes \mathcal{H}(P)$. Then

$$(F \otimes F)(\mathcal{R}) = \sum_{i,t=1}^{m} \langle \varepsilon_P, a_t \rangle b_t a_i \otimes b_i = \sum_{i,t=1}^{m} \langle b_t^*, 1_A \rangle b_t a_i \otimes b_i = \sum_{i=1}^{m} a_i \otimes b_i = R.$$

4) Consider the map

$$\rho = \rho_P : A \to P \otimes A, \quad \rho(a) = (1_A \otimes a)R = \sum R^1 \otimes aR^2 = \sum_{i=1}^{m} a_i \otimes ab_i$$

for all $a \in A$. We will show that $(A, \cdot, \rho_P) \in {}^P\mathcal{M}_P$ is a right-left P-Hopf module, where the structure of right P-module is simply the multiplication $\cdot$ of A. Indeed, for $a \in A$ we have

$$\begin{aligned}
(\mathrm{Id} \otimes \rho)\rho(a) &= \sum R^1 \otimes \rho(aR^2) = \sum R^1 \otimes r^1 \otimes aR^2 r^2 \\
&= (1_A \otimes 1_A \otimes a) R^{13} R^{23} \overset{(5)}{=} (1_A \otimes 1_A \otimes a)(R^{12})^{-1} R^{23} R^{12} \\
&= \sum U^1 r^1 \otimes U^2 R^1 r^2 \otimes aR^2 = \sum R^{-1}(1_A \otimes R^1) R \otimes aR^2 \\
&= \sum \Delta_P(R^1) \otimes aR^2 = (\Delta_P \otimes \mathrm{Id})\rho(a)
\end{aligned}$$

and

$$\sum_{i=1}^{m} \langle \varepsilon_P, a_i \rangle ab_i = \sum_{i=1}^{m} \langle b_i^*, 1_A \rangle ab_i = a$$

so (A,ρ) is a left P-comodule. The compatibility relation

$$\rho(a)\Delta_P(a_i) = (1_A \otimes a)RR^{-1}(1_A \otimes a_i)R = (1_A \otimes aa_i)R = \rho(aa_i)$$

holds for all $i = 1, \cdots, m$ and $a \in A$. Hence, $(A, \cdot, \rho_P) \in {}^P\mathcal{M}_P$ and the coinvariants

$$A^{\mathrm{co}(P)} = \{a \in A \mid \rho(a) = 1 \otimes a\} = A^{R,r}$$

are the right R-coinvariants of A. From the right-left version of the Fundamental Theorem of Hopf modules it follows that the multiplication of A,

$$\mu : A^{R,r} \otimes P \to A, \quad \mu(a \otimes x) = ax$$

defines an isomorphism of P-Hopf modules and, in particular, of right P-modules. We recall that $A^{R,r} \otimes P$ is a right P-module via $(a \otimes x) \cdot y = a \otimes xy$, for all $a \in A^{R,r}$, x, $y \in P$. It follows that A is free as a right P-module and, if A is finite dimensional,

$$\dim(A) = \dim(P)\dim(A^{R,r}).$$

In a similar way we can show that $(A, \cdot, \rho_H) \in {}_H\mathcal{M}^H$, where $\cdot$ is the multiplication of A and

$$\rho_H : A \to A \otimes H, \quad \rho_H(a) = R(a \otimes 1_A) = \sum R^1 a \otimes R^2$$

for all $a \in A$. Moreover, $A^{\mathrm{co}(H)} = A^{R,l}$. If we apply once again the Fundamental Theorem of Hopf modules (this time the left-right version) we obtain the other part of the statement.
5) $S = (f \otimes f)(R) = \sum_{i=1}^m f(a_i) \otimes f(b_i)$, $l(S) = l(R)$ and $S^{-1} = \sum f(U^1) \otimes f(U^2)$. It follows that $\{f(a_i) \mid i = 1, \cdots, m\}$ is a basis of $P(B,S)$ and hence the restriction of f to $P(A,R)$ gives an algebra isomorphism between $P(A,R)$ and $P(B,S)$ that is also a coalgebra map since

$$\begin{aligned} (f \otimes f)\Delta_{P(A,R)}(a_i) &= \sum f(U^1)f(R^1) \otimes f(U^2)f(a_i)f(R^2) \\ &= S^{-1}(1 \otimes f(a_i))S = \Delta_{P(B,S)}(f(a_i)) \end{aligned}$$

for all $i = 1, \cdots, m$. The last statement is obtain for $B = A$ and $f : A \to A$, $f(x) = uxu^{-1}$ for all $x \in A$. □

Corollary 2.2 *Let A be an algebra and $R \in A \otimes A$ an invertible solution of the pentagon equation. Then:*
1) R is unitary if and only if the subspaces $A^{R,l}$ and $A^{R,r}$ are nonzero.
2) If A is finite dimensional, then R is unitary.

Proof. 1. We have proved in Remark 1.4 that if the spaces $A^{R,l}$ and $A^{R,r}$ are nonzero, then R is unitary. Conversely, if R is unitary, the isomorphisms of vector spaces from part 4) of Theorem 2.1 show us that $A^{R,l}$ and $A^{R,r}$ are nonzero.

2. Assume now that A is finite dimensional. From the construction of Theorem 2.1 we have that $R \in P \otimes H$. In particular, for any positive integer t there exist scalars $\alpha_{ij} \in k$ such that

$$R^t = \sum_{i,j=1}^{m} \alpha_{ij} a_i \otimes b_j \tag{24}$$

We shall prove now that R is unitary, i.e. $1_A \in P \cap H$. As A is finite dimensional, A can be embedded into a matrix algebra $A \subset \mathcal{M}_n(k)$, where $n = \dim(A)$. We consider

$$R \in A \otimes A \subset \mathcal{M}_n(k) \otimes \mathcal{M}_n(k) \cong \mathcal{M}_{n^2}(k)$$

As R is invertible, using the Hamilton-Cayley theorem in $\mathcal{M}_{n^2}(k)$, the identity matrix I_{n^2} can be represented as a linear combination of powers of R. Hence, using (24), we obtain in $A \otimes A$ a linear combination $1_A \otimes 1_A = \sum_{i,j=1}^{m} \gamma_{ij} a_i \otimes b_j$, for some scalars $\gamma_{ij} \in k$. Therefore

$$1_A = \sum_{i,j=1}^{m} a_i \langle 1_A^*, \gamma_{ij} b_j \rangle = \sum_{i,j=1}^{m} \langle 1_A^*, \gamma_{ij} a_i \rangle b_j \in P \cap H$$

i.e. R is unitary. □

Let R be a unitary invertible solution of the pentagon equation. We have seen in Theorem 2.1 that the subalgebra A_R generated by the coefficient spaces $R_{(l)}$, $R_{(r)}$ is finite dimensional and, of course, we can view $R \in A_R \otimes A_R$. From this reason, in studying of the unitary invertible solutions of the pentagon equation, it is enough to focus on the case when the that the algebra A is finite dimensional. In this case, the unitary condition of R follows automatically.
Using Theorem 1.7 and Theorem 2.1 we obtain the following Corollary, which is the algebraic version of [2, Theorem 4.7]: the role of the operator V_S is played by the canonical element of a Heisenberg double.

Corollary 2.3 *$(A, R) \in$ Pent if and only if there exists a finite dimensional Hopf algebra L and an algebra map $F : \mathcal{H}(L) \to A$ such that $R = (F \otimes F)(\mathcal{R})$.*

Remark 2.4 Let $(A, R) \in$ *Pent*. It follows from Corollary 2.3 and Proposition 1.1 that there exists an algebra map $F : \mathcal{M}_m(k) \to A$, where $m = l(R)$; F is injective since $\mathcal{M}_m(k)$ is a simple algebra. Let $0 \neq a_{ij} = F(e_{ij}) \in A$, $i, j = 1, \cdots, m$; then $a_{ij} a_{kl} = \delta_{jk} a_{il}$ and $1_A = \sum_{i=1}^{m} a_{ii}$. It follows from the Reconstruction Theorem of the matrix algebra ([10, Theorem 17.5]) that there exists an algebra isomorphism

$$A \cong \mathcal{M}_m(B), \text{ where } B = \{x \in A \mid x a_{ij} = a_{ij} x, \ \forall i, j = 1, \cdots, m\}.$$

Hence A is a noncommutative algebra if R is non-trivial ($l(R) > 1$ or, equivalently, $R \neq 1_A \otimes 1_A$). Furthermore, $\dim(A) = m^2 \dim(B)$ and hence, $l(R)^2 | \dim(A)$.

We are now going to describe the space of integrals of the Hopf algebras $P = P(A, R)$ and $H = H(A, R)$, where $(A, R) \in \underline{Pent}$. Since A is free as a right P-module and as a left H-module, there exists

$$\pi_P : A \to P, \quad (\text{respectiv} \quad \pi_H : A \to H)$$

a non-zero right P-linear (respectiv left H-linear) map.

Proposition 2.5 *Let $(A, R) \in \underline{Pent}$. Then:*
1) $\pi_P(a)$ is a right integral in $P(A, R)$ for any right R-invariant $a \in {}^{R,r}A$.
2) $\pi_H(a)$ is a left integral in $H(A, R)$ for any left R-invariant $a \in {}^{R,l}A$.

Proof. 1. a is a right R-invariant; hence, $\sum_{i=1}^{m} aa_i \otimes b_i = a \otimes 1_A$. If we apply b_p^* to this equality we get that $aa_p = \langle b_p^*, 1_A \rangle a = \varepsilon_P(a_p)a$, for any $p = 1, \cdots, m$. As (a_i) is a basis of P, we obtain that $ax = \varepsilon_P(x)a$, for all $x \in P$. If we apply the right P-module map π_P to this, we obtain that $\pi_P(a)$ is a right integral of P.
2. Left to the reader. $\square$

3 The structure and the classification of finite dimensional Hopf algebras

We shall now prove the following Spliting Theorem:

Theorem 3.1 *Let L be a finite dimensional Hopf algebra. Then there exists an isomorphism of Hopf algebras*

$$L \cong P(\mathcal{H}(L), \mathcal{R})$$

where $\mathcal{R}$ is the canonical element of the Heisenberg double $\mathcal{H}(L)$.

Proof. Let $\{e_i, e_i^* \mid i = 1, \cdots, m\}$ be a basis of L and

$$\mathcal{R} = \sum_{i=1}^{m} (e_i \# \varepsilon_L) \otimes (1_L \# e_i^*) \in \mathcal{H}(L) \otimes \mathcal{H}(L)$$

the canonical element. We have to prove that the Hopf algebra $P(\mathcal{H}(L), \mathcal{R})$ extracted from part 1) of Theorem 2.1 is isomorphic to L, with the initial Hopf algebra structure. $i_L : L \to \mathcal{H}(L)$, $i_L(l) = l \# \varepsilon_L$ is an injective algebra map. We identify

$$L \cong \operatorname{Im}(i_L) = L \# \varepsilon_L.$$

From the construction, $P(\mathcal{H}(L), \mathcal{R})$ is the subalgebra of $\mathcal{H}(L)$ having $(e_i \# \varepsilon_L)_{i=1,\cdots,m}$ as a basis; i.e. there exists an algebra isomorphism $L \cong \operatorname{Im}(i_L) = P(\mathcal{H}(L), \mathcal{R})$. It remains to be proven that the coalgebra structure (resp. the antipode) of $P(\mathcal{H}(L), \mathcal{R})$ extracted from Theorem 2.1 is exactly the original coalgebra structure (resp. the antipode) of L. As the counit and the antipode of a Hopf algebra are uniquely determined by the multiplication and the comultiplication, the only thing left to be shown is the fact that, via the above identification, $\Delta_P = \Delta_L$. This means that

$$\Delta_L(e_i \# \varepsilon_L) = \mathcal{R}^{-1}(1_{\mathcal{H}(L)} \otimes e_i \# \varepsilon_L)\mathcal{R}$$

or equivalently,

$$\mathcal{R}\Delta_L(e_i \# \varepsilon_L) = \Big((1_L \# \varepsilon_L) \otimes (e_i \# \varepsilon_L)\Big)\mathcal{R}.$$

Now we compute

$$\begin{aligned} \Big((1_L \# \varepsilon_L) \otimes (e_i \# \varepsilon_L)\Big)\mathcal{R} &= \Big((1_L \# \varepsilon_L) \otimes (e_i \# \varepsilon_L)\Big)\Big(\sum_{j=1}^{m}(e_j \# \varepsilon_L) \otimes (1_L \# e_j^*)\Big) \\ &= \sum_{j=1}^{m}(e_j \# \varepsilon_L) \otimes (e_{i_{(2)}} \# e_{i_{(1)}} \cdot e_j^*) \end{aligned}$$

On the other hand

$$\begin{aligned} \mathcal{R}\Delta_L(e_i \# \varepsilon_L) &= \sum_{j=1}^{m}\Big((e_j \# \varepsilon_L) \otimes (1_L \# e_j^*)\Big)\Big((e_{i_{(1)}} \# \varepsilon_L) \otimes (e_{i_{(2)}} \# \varepsilon_L)\Big) \\ &= \sum_{j=1}^{m}(e_j e_{i_{(1)}} \# \varepsilon_L) \otimes (e_{i_{(2)}} \# e_j^*) \end{aligned}$$

Hence, we have to show the formula

$$\sum_{j=1}^{m} e_j e_{i_{(1)}} \otimes e_{i_{(2)}} \otimes e_j^* = \sum_{j=1}^{m} e_j \otimes e_{i_{(2)}} \otimes e_{i_{(1)}} \cdot e_j^* \tag{25}$$

Let us set the indices a, b, $k = 1, \cdots, m$, and evaluate (25) at $e_a^* \otimes e_b^* \otimes e_k$. (25) is then equivalent to

$$\langle e_a^*, e_k e_{i_{(1)}}\rangle\langle e_b^*, e_{i_{(2)}}\rangle = \sum_{j=1}^{m}\langle e_a^*, e_j\rangle\langle e_b^*, e_{i_{(2)}}\rangle\langle e_{i_{(1)}} \cdot e_j^*, e_k\rangle$$

and this is easily verified using the dual basis formula. It follows that $\Delta_L = \Delta_P$ and $L \cong P(\mathcal{H}(L), \mathcal{R})$ as Hopf algebras. $\square$

We have now arrived at the Structure Theorem for finite dimensional Hopf algebras. Let L be a finite dimensional Hopf algebra. Proposition 1.1 proves that there exists an algebra isomorphism $\mathcal{H}(L) \cong \mathcal{M}_n(k)$, where $n = \dim(L)$. Via this isomorphism the canonical element $\mathcal{R} \in \mathcal{H}(L) \otimes \mathcal{H}(L)$ is viewed as an element of $\mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$, or as a matrix of $\mathcal{M}_{n^2}(k)$. We shall now give the data which shows us how any finite dimensional Hopf algebra is constructed. Let $R \in \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$ be an invertible solution of the pentagon equation. We write $R = \sum_{i=1}^{m} A_i \otimes B_i$ where $m = l(R)$; then the sets of matrices $\{A_i \mid i = 1, \cdots m\}$ and $\{B_i \mid i = 1, \cdots m\}$ are linearly independent over k. Let $\{B_i^* \mid i = 1, \cdots m\}$ be the dual basis of $\{B_i \mid i = 1, \cdots m\}$ and $U = R^{-1} = \sum U^1 \otimes U^1$.

The Hopf algebra $P(\mathcal{M}_n(k), R)$ will be denoted for simplicity by $P(n, R)$ and is described as follows:

- as an algebra, $P(n, R)$ is the subalgebra of the $n \times n$-matrix algebra $\mathcal{M}_n(k)$ with $\{A_i \mid i = 1, \cdots m\}$ as a k-basis;
- the coalgebra structure and the antipode of $P(n, R)$ are given by the following formulas:

$$\Delta : P(n, R) \to P(n, R) \otimes P(n, R), \quad \Delta(A_p) = R^{-1}(I_n \otimes A_p)R \tag{26}$$

$$\varepsilon : P(n, R) \to k, \quad \varepsilon(A_p) = \langle B_p^*, I_n \rangle \tag{27}$$

$$S : P(n, R) \to P(n, R), \quad S(A_p) = \sum \langle B_p^*, U^2 \rangle U^1 \tag{28}$$

for all $p = 1, \cdots, m$. The dual of $P(n, R)$ is the subalgebra $H(n, R) = H(\mathcal{M}_n(k), R)$ of the matrix algebra $\mathcal{M}_n(k)$ having $\{B_i \mid i = 1, \cdots m\}$ as a k-basis and the comultiplication given by

$$\Delta_l : H(n, R) \to H(n, R) \otimes H(n, R), \quad \Delta_l(B_p) = R(B_p \otimes I_n)R^{-1}$$

for all $p = 1, \cdots, m$.
Theorems 2.1 and 3.1 imply the following Structure Theorem for finite dimensional Hopf algebras.

Theorem 3.2 *L is a finite dimensional Hopf algebra if and only if there exists a positive integer n and an invertible solution of the pentagon equation $R \in \mathcal{M}_n(k) \otimes \mathcal{M}_n(k) \cong \mathcal{M}_{n^2}(k)$ such that $L \cong P(n, R)$. Furthermore,*

$$\dim(L) = \frac{n^2}{\dim(\mathcal{M}_n(k)^{R,r})} = l(R). \tag{29}$$

Let n be a positive integer. The algebra isomorphism $\mathcal{H}(L) \cong \mathcal{M}_{\dim(L)}(k)$ and Theorem 3.1 show us that an n-dimensional Hopf algebra L is isomorphic to a $P(n, R)$ for $R \in \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$ an invertible solution of the pentagon equation such that $l(R) = n$. Overall, each solution of the pentagon equation belonging to $\mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$ determines a Hopf algebra, the dimension of which is not necessarily n, but a divisor of n^2 based on the Lagrange formula (29).
We are now going to prove the Classification Theorem for finite dimensional Hopf algebras. Let $\underline{Pent}_n$ be the set

$$\underline{Pent}_n = \{R \in \mathcal{M}_n(k) \otimes \mathcal{M}_n(k) \ \mid \ (\mathcal{M}_n(k), R) \in \underline{Pent} \text{ and } l(R) = n\}.$$

Theorem 3.3 *Let n be a positive integer. Then there exists a one to one correspondence between the set of types of n-dimensional Hopf algebras and the set of the orbits of the action*

$$GL_n(k) \times \underline{Pent}_n \to \underline{Pent}_n, \quad (u, R) \to (u \otimes u)R(u \otimes u)^{-1}. \tag{30}$$

Proof. In part 5) of the Theorem 2.1 we proved that there exists a Hopf algebra isomorphism $P(n,R) \cong P(n, {}^uR)$ for any $R \in \underline{Pent}_n$ and $u \in GL_n(k)$, which means that all the Hopf algebras associated to the elements of an orbit of the action (30) are isomorphic. We shall now prove the converse. First we show that two finite dimensional Hopf algebras L_1 and L_2 are isomorphic if and only if $(\mathcal{H}(L_1), \mathcal{R}_{L_1})$ and $(\mathcal{H}(L_2), \mathcal{R}_{L_2})$ are isomorphic as objects in $\underline{Pent}$. The "only if" part follows from Theorem 3.1 and part 5) of Theorem 2.1. It remains to prove the "if" part. Let $f : L_1 \to L_2$ be a Hopf algebra isomorphism. Then, $f^* : L_2^* \to L_1^*$, $f^*(l^*) = l^* \circ f$ is an isomorphism of Hopf algebras and

$$\tilde{f} : \mathcal{H}(L_1) \to \mathcal{H}(L_2), \quad \tilde{f}(h\#h^*) := f(h)\#(f^*)^{-1}(h^*) = f(h)\#h^* \circ f^{-1}$$

for all $h \in L_1$, $h^* \in L_1^*$ is an algebra isomorphism between the two Heisenbergs. On the other hand, if $\{e_i, e_i^*\}$ is a dual basis of L_1, then $\{f(e_i), e_i^* \circ f^{-1}\}$ is a dual basis of L_2 and hence $(\tilde{f} \otimes \tilde{f})(\mathcal{R}_{L_1}) = \mathcal{R}_{L_2}$, and this proves that $\tilde{f}$ is an isomorphism in $\underline{Pent}$. Let $n_i = \dim(L_i)$, $i = 1, 2$. Using Proposition 1.1 we obtain that $(\mathcal{H}(L_1), \mathcal{R}_{L_1}) \cong (\mathcal{H}(L_2), \mathcal{R}_{L_2})$ if and only if $(\mathcal{M}_{n_1}(k), R_1) \cong (\mathcal{M}_{n_2}(k), R_2)$ in $\underline{Pent}$, where R_i is the image of $\mathcal{R}_{L_i}$ under the algebra isomorphism $\mathcal{H}(L_i) \cong \mathcal{M}_{n_i}(k)$. Now, the two matrix algebras $\mathcal{M}_{n_1}(k)$ and $\mathcal{M}_{n_2}(k)$ are isomorphic if and only if $n_1 = n_2$ and the Skolem-Noether theorem tells us that any automorphism g of the matrix algebra $\mathcal{M}_{n_1}(k)$ is an inner one: there exists $u \in GL_{n_1}(k)$ such that $g(x) = g_u(x) = uxu^{-1}$. Hence we obtain that $(\mathcal{M}_{n_1}(k), R_1) \cong (\mathcal{M}_{n_2}(k), R_2)$ in $\underline{Pent}$ if and only if $n_1 = n_2$ and there exists $u \in GL_{n_1}(k)$ such that $R_2 = (g_u \otimes g_u)(R_1) = (u \otimes u)R_1(u \otimes u)^{-1}$, i.e. R_2 is equivalent to R_1, as needed. □

We shall conclude with a few examples, evidencing first of all the general method of determining invertible solutions of the pentagon equation $R \in \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$. Let $(e_{ij})_{i,j=1,n}$ be the canonical basis of $\mathcal{M}_n(k)$. An element $R \in \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$ can be written as follows

$$R = \sum_{i,j=1}^{n} e_{ij} \otimes A_{ij} \tag{31}$$

for some matrices $A_{ij} \in \mathcal{M}_n(k)$. Using the formula (1), R viewed as a matrix in $\mathcal{M}_{n^2}(k)$, is given by $R = (A_{ij})_{i,j=1,\cdots,n}$, and we can quichly check if R is invertible ($\det(R) \neq 0$). Let us clarify the condition for R to be a solution of the pentagon equation. Taking into account the multiplication rule among the elements (e_{ij}) we have:

$$R^{12}R^{13}R^{23} = \sum_{i,j,p,r,s=1}^{n} e_{ip} \otimes A_{ij}e_{rs} \otimes A_{jp}A_{rs} \quad \text{and} \quad R^{23}R^{12} = \sum_{a,b,i,p=1}^{n} e_{ip} \otimes e_{ab}A_{ip} \otimes A_{ab}.$$

Hence, R is a solution of the pentagon equation if and only if

$$\sum_{a,b=1}^{n} e_{ab}A_{ip} \otimes A_{ab} = \sum_{i,r,s=1}^{n} A_{ij}e_{rs} \otimes A_{jp}A_{rs} \tag{32}$$

or equivalently,

$$R(A_{ip} \otimes I_n) = (\sum_{j=1}^{n} A_{ij} \otimes A_{jp})R \tag{33}$$

for all $i,\ p = 1, \cdots n$, which can be viewed as an equation in $\mathcal{M}_{n^2}(k)$. We record this observation in the following:

Proposition 3.4 *Let n be a positive integer and $R = (A_{ij})_{i,j=1,n} \in \mathcal{M}_{n^2}(k) \cong \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$, an invertible matrix. Then R is a solution of the pentagon equation if and only if*

$$\sum_{j=1}^{n} A_{ij} \otimes A_{jp} = R(A_{ip} \otimes I_n)R^{-1} \tag{34}$$

for all $i,\ p = 1, \cdots, n$.

Examples 3.5 1. Let n be a positive integer, $A = e_{21} + e_{32} + \cdots + e_{n,n-1} + e_{1n} \in \mathcal{M}_n(k)$ and $A_{ij} = \delta_{ij} A^{i-1}$ for all $i,\ j = 1, \cdots n$. Then

$$R = e_{11} \otimes I_n + e_{22} \otimes A + \cdots + e_{nn} \otimes A^{n-1} \in \mathcal{M}_n(k) \otimes \mathcal{M}_n(k)$$

is an invertible solution of the pentagon equation and $P(n, R) \cong (kG)^*$, the Hopf algebra of functions on a cyclic group with n elements G. Indeed, the pentagon equation (33) becomes

$$R(A_{ii} \otimes I_n) = (A_{ii} \otimes A_{ii})R$$

for all $i = 1, \cdots, n$. As $A_{ii} = A^{i-1}$, it is enough if we prove that $R(A \otimes I_n) = (A \otimes A)R$. Using the expression of A and the fact that $A^n = I_n$, we have

$$R(A \otimes I_n) = (A \otimes A)R = e_{1n} \otimes I_n + e_{21} \otimes A + \cdots + e_{n,n-1} \otimes A^{n-1}$$

i.e. (33) holds and R is an invertible solution of the pentagon equation. It remains to prove that $P(n, R) \cong (kG)^*$. We shall prove that $H = H(n, R) \cong kG$, the group algebra of G and then use the duality between $P(n, R)$ and $H(n, R)$ given by Theorem 2.1. We remark that R is already written in the form $R = \sum_{i=1}^{n} A_i \otimes B_i$, where (A_i) and (B_i) are linearly independent. Then $H(n, R)$ is the commutative subalgebra of $\mathcal{M}_n(k)$ with $\{I_n, A, A^2, \cdots, A^{n-1}\}$ as a basis. Using (18), (22) and the fact that $R^{-1} = e_{11} \otimes I_n + e_{22} \otimes A^{n-1} + \cdots + e_{nn} \otimes A$, we obtain that the comultiplication, the counit and the antipode of H are given by

$$\Delta_H(A) = A \otimes A, \quad \varepsilon_H(A) = I_n, \quad S_H(A) = A^{n-1} = A^{-1}$$

i.e. $H \cong kG$.
2. Let $R \in \mathcal{M}_4(k) \otimes \mathcal{M}_4(k)$ given by

$$R = (e_{11} + e_{44}) \otimes I_4 + (e_{22} + e_{33}) \otimes (e_{12} + e_{21} + e_{34} + e_{43}) + e_{13} \otimes (e_{31} - e_{42}) + e_{24} \otimes (e_{41} - e_{32})$$

Then R is an invertible solution of the pentagon equation $H(4, R) \cong H_4$, and hence $P(4, R) \cong H_4^* \cong H_4$, where H_4 is the Sweedler four dimensional noncommutative noncocommutative Hopf algebra.
The proof is similar to the previous example (we left to the reader to check the pentagon equation (33) for R). The inverse of R is

$$R^{-1} = (e_{11} + e_{44}) \otimes I_4 + (e_{22} + e_{33}) \otimes (e_{12} + e_{21} + e_{34} + e_{43}) + e_{13} \otimes (e_{41} - e_{32}) + e_{24} \otimes (e_{24} - e_{31}).$$

and therefore the Hopf algebra $H(4, R)$ is the four dimensional subalgebra of $\mathcal{M}_4(k)$ having

$$\{I_4, e_{12} + e_{21} + e_{34} + e_{43}, e_{41} - e_{32}, e_{31} - e_{42}\}$$

as a k-basis. Now, writing $x = e_{31} - e_{42}$ and $g = e_{12} + e_{21} + e_{34} + e_{43}$ we find that

$$x^2 = 0, \quad g^2 = I_4, \quad gx = -xg = e_{41} - e_{32}$$

On the other hand, the formula of the comultiplication of $H(4, R)$ given by (18), namely $\Delta(A) = R(A \otimes I_4)R^{-1}$ for all $A \in H(4, R)$, gives, using the expression of R^{-1},

$$\Delta(g) = g \otimes g, \quad \Delta(x) = x \otimes g + I_4 \otimes x$$

i.e. $H(4, R) \cong H_4$, the Sweedler four dimensional Hopf algebra.

4 Conclusions and outlooks

The study of the pentagon equation, together with a few classic results in algebra (the Fundamental Theorem for Hopf modules, Morita theory and the Skolem-Noether theorem), have led us to the Structure and Classification theorem for finite dimensional Hopf algebras. Theorem 3.3 opens a new road for describing the types of isomorphisms for Hopf algebras of a certain dimension. The first step, and the most important, is however the development of a new Jordan type theory (we called it *restricted Jordan theory*). From the point of view of actions, the classical Jordan theory gives the most elementary description of the representatives of the orbits of the action

$$GL_n(k) \times \mathcal{M}_n(k) \to \mathcal{M}_n(k), \quad (U, A) \to UAU^{-1}.$$

The restricted Jordan theory refers to the following open problem:
Problem: Describe the orbits of the action

$$GL_n(k) \times (\mathcal{M}_n(k) \otimes \mathcal{M}_n(k)) \to \mathcal{M}_n(k) \otimes \mathcal{M}_n(k), \quad (U, R) \to (U \otimes U)R(U \otimes U)^{-1}.$$

We recall that the canonical Jordan form J_A of a matrix A is the matrix equivalent to A which has the greatest number of zeros. For practical reasons, in the restricted Jordan theory we are in fact interested in finding the elements of each orbit that have the greatest number of zeros. Of these, we retain only those which are invertible solutions of the pentagon equation. The set of types of n-dimensional Hopf algebras shall be those Hopf algebras associated (using Theorem 3.2) to the solutions of length n (or, equivalently, the space of coinvariant elements is n-dimensional); all other Hopf algebras will have a dimension that is a divisor of n^2. We mention that, as a general rule, the set of types of n-dimensional Hopf algebras is infinite (this was proved recently in [1], [3], [7]). If however we limit ourselves to classifying certain special types of Hopf algebras, then this set can be finite. For instance, the set of types of n-dimensional semisimple and cosemisimple Hopf algebras is finite ([14]).

References

[1] N. Andruskiewitsch and H.-J. Schneider, Lifting of quantum linear space and pointed Hopf algebras of order p^3, *J. Algebra* **209** (1998), 658-691.

[2] S. Baaj and G. Skandalis, Unitaries multiplicatifs et dualite pour les produits croises de C^*-algebres, *Ann. Sci. Ecole Norm. Sup.* **26** (1993), 425–488.

[3] M. Beattie, S. Dascalescu and L. Grunenfelder, On the number of types of finite dimensional Hopf algebrasm, *Inv. Math.* **136** (1999), 1-7.

[4] S. Caenepeel, Bogdan Ion, G. Militaru and S. Zhu, Separable functors for Doi-Hopf modules. Applications, *Advances Math.* **145** (1999), 239-290.

[5] A. A. Davydov, Pentagon equation and matrix bialgebras, q-alg/0001095.

[6] Y. Doi, Unifying Hopf modules, *J. Algebra* **153** (1992), 373–385.

[7] S. Gelaki, On pointed Hopf algebras and Kaplansky's 10th conjecture, *J. Algebra* **209** (1998), 635-657.

[8] R.M. Kashaev, Heisenberg Double and the Pentagon relation, *Algebra i Analiz* **8** (1996), 63-74; translation in *St. Petersburg Math. J.* **8** (1997), 585-592; q-alg/9503005.

[9] R.M. Kashaev and S.M. Sergeev, On pentagon, ten-term and tetraedron relations, *Comm. Math. Phys.* **195** (1998), 309-319.

[10] T.Y Lam, Lectures on modules and rings, Springer-Verlag, 1998.

[11] Di-Ming Lu, Tannaka duality and the FRT-construction, *Comm. Algebra*, to appear.

[12] G. Militaru, The Hopf modules category and the Hopf equation, *Comm. Algebra* **10** (1998), 3071-3097.

[13] S. Montgomery, Classifying finite-dimensional semisimple Hopf algebras, *AMS Contemp. Math.* **229** (1998), 265-279.

[14] D. Stefan, The set of types of n-dimensional semisimple and cosemisimple Hopf algebras is finite, *J. Algebra* **193** (1997), 571-580.

[15] R. Street, Fusion operators and cocycloids in monoidal categories, *Appl. Categorical Structures*, **6** (1998), 177-191.

[16] M. E. Sweedler, "Hopf algebras", Benjamin, New York, 1969.

[17] A. Van Daele and S. Van Keer, The Yang-Baxter and Pentagon equation, *Compositio Math.* **91** (1994), 201-221.